

THE GROUP OF CAUSAL AUTOMORPHISMS

DO-HYUNG KIM

ABSTRACT. The group of causal automorphisms on Minkowski space-time is given and its structure is analyzed.

1. INTRODUCTION

In 1964, Zeeman has shown that any causal automorphism on Minkowski space-time $\mathbb{R}_1^n$ can be represented by a composite of orthochronous transformation, traslation and homothety when $n \geq 3$. (See Ref. [1]). Therefore, if we let G be the group of all causal automorphisms on $\mathbb{R}_1^n$ with $n \geq 3$, then G is isomorphic to the semi-direct product of $\mathbb{R}^+ \times \mathcal{O}$ and $\mathbb{R}_1^n$ where $\mathbb{R}^+$ is the group of positive real numbers and $\mathcal{O}$ is the group of orthochronous transformations on $\mathbb{R}_1^n$. However, as he remarked, Zeeman's theorem does not hold when $n = 2$.

In connection with this, recently, the standard form of causal automorphism on $\mathbb{R}_1^2$ is given in Ref. [2]. In this paper, we improve the result in Ref. [2] and by use of this, the structure of the causal automorphism group is analyzed when $n = 2$.

2. CAUSAL AUTOMORPHISMS ON $\mathbb{R}_1^2$

The following is a known result.

Theorem 2.1. *Let $F : \mathbb{R}_1^2 \rightarrow \mathbb{R}_1^2$ be a causal automorphism. Then, there exist a continuous functions $g : \mathbb{R} \rightarrow \mathbb{R}$ and a homeomorphism $f : \mathbb{R} \rightarrow \mathbb{R}$ which satisfy $\sup(g \pm f) = \infty$, $\inf(g \pm f) = -\infty$ and $|\frac{g(t+\delta t)-g(t)}{f(t+\delta t)-f(t)}| < 1$ for all t and δt , such that if f is increasing, then F is given by*

$$F(x, y) = (\frac{f(x-y)+f(x+y)}{2} + \frac{g(x+y)-g(x-y)}{2}, \frac{f(x+y)-f(x-y)}{2} + \frac{g(x-y)+g(x+y)}{2})$$

and if f is decreasing, then we have

$$F(x, y) = (\frac{f(x+y)+f(x-y)}{2} + \frac{g(x-y)-g(x+y)}{2}, \frac{f(x-y)-f(x+y)}{2} + \frac{g(x+y)+g(x-y)}{2}).$$

Conversely, for any functions f and g which satisfy the above conditions, the function $F : \mathbb{R}_1^2 \rightarrow \mathbb{R}_1^2$ defined as above, is a causal automorphism.

Proof. See Theorem 4.4 in Ref. [2]. □

Key words and phrases. Lorentzian geometry, general relativity, causality, Cauchy surface, space-time, global hyperbolicity, causal automorphism, Minkowski space-time.

We remark that in the above theorem, for given causal automorphism $F(x, y) = (F_1(x, y), F_2(x, y))$, the homeomorphism f and the continuous function g are uniquely determined by $f(t) = F_1(t, 0)$ and $g(t) = F_2(t, 0)$. By use of this, we improve the above theorem.

Theorem 2.2. *Let $F : \mathbb{R}_1^2 \rightarrow \mathbb{R}_1^2$ be a causal automorphism on $\mathbb{R}_1^2$. Then, there exist unique homeomorphisms φ and ψ of $\mathbb{R}$, which are either both increasing or both decreasing, such that if φ and ψ are increasing, then we have $F(x, y) = (\frac{1}{2}(\varphi(x+y) + \psi(x-y)), \varphi(x+y) - \psi(x-y))$, or if φ and ψ are decreasing, then we have $F(x, y) = (\frac{1}{2}(\varphi(x-y) + \psi(x+y)), \varphi(x-y) - \psi(x+y))$.*

Conversely, for any given homeomorphisms φ and ψ of $\mathbb{R}$, which are either both increasing or both decreasing, the function F defined as above is a causal automorphism of $\mathbb{R}_1^2$.

Proof. For any given causal automorphism $F : \mathbb{R}_1^2 \rightarrow \mathbb{R}_1^2$, by the previous theorem, we can get unique homeomorphism f and unique continuous function g . Let $\varphi = f + g$ and $\psi = f - g$. Then, clearly, φ and ψ are continuous.

The conditions $\sup(g \pm f) = \infty$, $\inf(g \pm f) = -\infty$ imply that φ and ψ are surjective. If $\varphi(t) = \varphi(t')$ with $t \neq t'$, then this implies that $\frac{g(t') - g(t)}{f(t') - f(t)} = -1$, which contradicts to the condition $|\frac{g(t+\delta t) - g(t)}{f(t+\delta t) - f(t)}| < 1$ for all t and δt . Therefore, φ is injective and likewise, we can show that ψ is injective. Since φ and ψ are continuous bijections from $\mathbb{R}$ to $\mathbb{R}$, the topological domain of invariance implies that φ and ψ are homeomorphisms.

The equation $\varphi(t_2) - \varphi(t_1) = (f(t_2) - f(t_1))[1 + \frac{g(t_2) - g(t_1)}{f(t_2) - f(t_1)}]$ tells us that, by the condition $|\frac{g(t+\delta t) - g(t)}{f(t+\delta t) - f(t)}| < 1$, φ is increasing if and only if f is increasing. Likewise, we can show that ψ is increasing if and only if f is increasing.

By simple calculation, we can show that F has the desired form when expressed in terms of φ and ψ . This completes the proof of the first part.

To prove the converse, let φ and ψ be increasing homeomorphisms on $\mathbb{R}$. Let $f = \frac{1}{2}(\varphi + \psi)$ and $g = \frac{1}{2}(\varphi - \psi)$. Then g is continuous and f is an increasing homeomorphism. Since φ and ψ are homeomorphisms, we have $\sup(f + g) = \sup \varphi = \infty$, $\sup(f - g) = \sup \psi = \infty$, $\inf(f + g) = \inf \varphi = -\infty$ and $\inf(f - g) = \inf \psi = -\infty$.

We now show that f and g satisfy the inequality $|\frac{g(t+\delta t) - g(t)}{f(t+\delta t) - f(t)}| < 1$. Let $\Delta = \frac{g(t) - g(t_0)}{f(t) - f(t_0)} = \frac{\varphi(t) - \varphi(t_0) - (\psi(t) - \psi(t_0))}{\varphi(t) - \varphi(t_0) + \psi(t) - \psi(t_0)}$. Without loss of generality, we can assume $t > t_0$ and it is easy to see that $-1 < \Delta < 1$, since φ and ψ are increasing. Therefore, by the theorem 2.1, the function F defined as in the statement, is a causal automorphism. By the exactly same argument, we can show that the assertion also holds when φ and ψ are both decreasing homeomorphisms. $\square$

3. THE GROUP OF CAUSAL AUTOMORPHISMS ON $\mathbb{R}_1^2$

In this section, we denote the group of all causal automorphisms on $\mathbb{R}_1^2$ by G , and we analyze its group structure. For this we let $H(\mathbb{R})$ be the group of all homeomorphisms on $\mathbb{R}$ and let $H = H^+ \cup H^-$ where $H^+ = \{(\varphi, \psi) \in H(\mathbb{R}) \times H(\mathbb{R}) \mid \varphi, \psi \text{ are increasing}\}$ and $H^- = \{(\varphi, \psi) \in H(\mathbb{R}) \times H(\mathbb{R}) \mid \varphi, \psi \text{ are decreasing}\}$. Then H is a subgroup of $H(\mathbb{R}) \times H(\mathbb{R})$ under the operation induced from $H(\mathbb{R}) \times H(\mathbb{R})$.

From the Theorem 2.2, we can see that any causal automorphism F on $\mathbb{R}_1^2$ corresponds to a unique element in H and, conversely, each elements in H uniquely determines a causal automorphism on $\mathbb{R}_1^2$. Thus, there exists a one-to-one correspondence between G and H as a set. It might seem that G is isomorphic to H . However, we cannot obtain an isomorphism in this way and we define a new operation on H as follows.

We define a $\mathbb{Z}_2$ -action on H by $a \cdot (\varphi, \psi) = (\varphi, \psi)$ if $a = 0$ and $a \cdot (\varphi, \psi) = (\psi, \varphi)$ if $a = 1$. If we define a map $\pi : H \rightarrow \mathbb{Z}_2$ by $\pi(x) = 0$ if $x \in H^+$ and $\pi(x) = 1$ if $x \in H^-$, then π is a group homomorphism with H equipped with the operation induced from $H(\mathbb{R}) \times H(\mathbb{R})$. Note that $\pi(\varphi, \psi) = \pi(\varphi^{-1}, \psi^{-1}) = \pi(\psi, \varphi)$ and $\pi(\pi(a, b) \cdot (\varphi, \psi)) = \pi(\varphi, \psi)$ for any $(a, b) \in H$.

To get an isomorphism from G to H , we define a new operation $*$ on H by $(\alpha, \beta) * (\varphi, \psi) = (\alpha, \beta) \circ \pi(\alpha, \beta) \cdot (\varphi, \psi)$ where $\cdot$ is the $\mathbb{Z}_2$ -action defined above and $\circ$ is the operation induced from $H(\mathbb{R}) \times H(\mathbb{R})$.

Theorem 3.1. *The set H under $*$ is a group and is isomorphic to G .*

Proof. To show associativity, by calculation, we have, $(a, b) * \{(c, d) * (e, f)\} = (a, b) * \{(c, d) \circ \pi(c, d) \cdot (e, f)\} = (a, b) \circ \pi(a, b) \cdot \{(c, d) \circ \pi(c, d) \cdot (e, f)\}$. Thus, we have the following four cases.

$\pi(a, b)$	$\pi(c, d)$	$(a, b) * \{(c, d) * (e, f)\}$
0	0	$(a, b) \circ \{(c, d) \circ (e, f)\} = (ace, bdf)$
0	1	$(a, b) \circ \{(c, d) \circ (f, e)\} = (acf, bde)$
1	0	$(a, b) \circ (df, ce) = (adf, bce)$
1	1	$(a, b) \circ (de, cf) = (ade, bcf)$

On the other hand, we have

$$\begin{aligned}
 \{(a, b) * (c, d)\} * (e, f) &= \{(a, b) \circ \pi(a, b) \cdot (c, d)\} * (e, f) \\
 &= \{(a, b) \circ \pi(a, b) \cdot (c, d)\} \circ \pi((a, b) \circ \pi(a, b) \cdot (c, d)) \cdot (e, f) \\
 &= \{(a, b) \circ \pi(a, b) \cdot (c, d)\} \circ \{\pi(a, b) \pi(\pi(a, b) \cdot (c, d))\} \cdot (e, f)
 \end{aligned}$$

($\cdot$: π is a group homomorphism.)

$$= \{(a, b) \circ \pi(a, b) \cdot (c, d)\} \circ \{\pi(a, b)\pi(c, d)\} \cdot (e, f)$$

$$(\cdot \pi(\pi(a, b) \cdot (c, d)) = \pi(c, d))$$

Thus, we have the following four cases.

$\pi(a, b)$	$\pi(c, d)$	$\{(a, b) * (c, d)\} * (e, f)$
0	0	$\{(a, b) \circ (c, d)\} \circ (e, f) = (ace, bdf)$
0	1	$\{(a, b) \circ (c, d)\} \circ (f, e) = (acf, bde)$
1	0	$\{(a, b) \circ (d, c)\} \circ (f, e) = (adf, bce)$
1	1	$\{(a, b) \circ (d, c)\} \circ (e, f) = (ade, bcf)$

The above two tables tell us that the operation $*$ satisfies the associative law.

The following two formulae show that (id, id) is the identity element in H under $*$.

$$\begin{aligned} (id, id) * (\varphi, \psi) &= (id, id) \circ \pi(id, id) \cdot (\varphi, \psi) \\ &= (id, id) \circ (\varphi, \psi) = (\varphi, \psi), \\ (\varphi, \psi) * (id, id) &= (\varphi, \psi) \circ \pi(\varphi, \psi) \cdot (id, id) \\ &= (\varphi, \psi) \circ (id, id) = (\varphi, \psi). \end{aligned}$$

The following two formulae show that, for given $(\varphi, \psi) \in H$, $\pi(\varphi, \psi) \cdot (\varphi^{-1}, \psi^{-1})$ is the inverse element of (φ, ψ) .

$$\begin{aligned} (\varphi, \psi) * \{\pi(\varphi, \psi) \cdot (\varphi^{-1}, \psi^{-1})\} &= (\varphi, \psi) \circ \pi(\varphi, \psi) \cdot \{\pi(\varphi, \psi) \cdot (\varphi^{-1}, \psi^{-1})\} \\ &= (\varphi, \psi) \circ \{\pi(\varphi, \psi)\pi(\varphi, \psi)\} \cdot (\varphi^{-1}, \psi^{-1}) \\ &= (\varphi, \psi) \circ (\varphi^{-1}, \psi^{-1}) = (id, id), \\ \{\pi(\varphi, \psi) \cdot (\varphi^{-1}, \psi^{-1})\} * (\varphi, \psi) &= \{\pi(\varphi, \psi) \cdot (\varphi^{-1}, \psi^{-1})\} \circ \{\pi(\pi(\varphi, \psi) \cdot (\varphi^{-1}, \psi^{-1}))\} \cdot (\varphi, \psi) \\ &= \{\pi(\varphi, \psi) \cdot (\varphi^{-1}, \psi^{-1})\} \circ \{\pi(\varphi, \psi) \cdot (\varphi, \psi)\} = (id, id). \end{aligned}$$

Therefore, H forms a group under the operation $*$.

We now show that the groups G and H under $*$ are isomorphic. For given causal automorphism $F(x, y) = (F_1(x, y), F_2(x, y))$, Theorem 2.2 tells us that $F_1 + F_2$ and $F_1 - F_2$ determine unique element in H . Therefore, we can define $\Pi : G \rightarrow H$ by

$$\Pi(F) = (\text{homeomorphism determined by } F_1 + F_2, \text{homeomorphism determined by } F_1 - F_2)$$

By Theorem 2.2, Π is a bijection and we only need to show that Π is a homomorphism. We have the following four cases.

i) Let $G = \frac{1}{2}(\alpha(x+y)+\beta(x-y), \alpha(x+y)-\beta(x-y))$ and $F = \frac{1}{2}(\varphi(x+y)+\psi(x-y), \varphi(x+y)-\psi(x-y))$ where both (α, β) and (φ, ψ) are in H^+ . Then, we have $G \circ F(x, y) = \frac{1}{2}(\alpha \circ \varphi(x+y) + \beta \circ \psi(x-y), \alpha \circ \varphi(x+y) - \beta \circ \psi(x-y))$ and thus $\Pi(G \circ F) = (\alpha \circ \varphi, \beta \circ \psi)$. Since $\Pi(G) = (\alpha, \beta)$ and $\Pi(F) = (\varphi, \psi)$, we have $\Pi(G \circ F) = \Pi(G) * \Pi(F)$.

ii) Let $G = \frac{1}{2}(\alpha(x+y) + \beta(x-y), \alpha(x+y) - \beta(x-y))$ and $F = \frac{1}{2}(\varphi(x-y) + \psi(x+y), \varphi(x-y) - \psi(x+y))$ where (α, β) is in H^+ and (φ, ψ) is in H^- . Then, we have $G \circ F(x, y) = \frac{1}{2}(\alpha \circ \varphi(x-y) + \beta \circ \psi(x+y), \alpha \circ \varphi(x-y) - \beta \circ \psi(x+y))$ and thus $\Pi(G \circ F) = (\alpha \circ \varphi, \beta \circ \psi)$. Since $\Pi(G) = (\alpha, \beta)$ and $\Pi(F) = (\varphi, \psi)$, we have $\Pi(G \circ F) = \Pi(G) * \Pi(F)$.

iii) Let $G = \frac{1}{2}(\alpha(x-y) + \beta(x+y), \alpha(x-y) - \beta(x+y))$ and $F = \frac{1}{2}(\varphi(x+y) + \psi(x-y), \varphi(x+y) - \psi(x-y))$ where (α, β) is in H^- and (φ, ψ) is in H^+ . Then, we have $G \circ F(x, y) = \frac{1}{2}(\alpha \circ \psi(x-y) + \beta \circ \varphi(x+y), \alpha \circ \psi(x-y) - \beta \circ \varphi(x+y))$ and thus $\Pi(G \circ F) = (\alpha \circ \psi, \beta \circ \varphi)$. Since $\Pi(G) = (\alpha, \beta)$ and $\Pi(F) = (\varphi, \psi)$, we have $\Pi(G \circ F) = (\alpha \circ \psi, \beta \circ \varphi) = (\alpha, \beta) \circ \pi(\alpha, \beta) \cdot (\varphi, \psi) = \Pi(G) * \Pi(F)$.

iv) Let $G = \frac{1}{2}(\alpha(x-y)+\beta(x+y), \alpha(x-y)-\beta(x+y))$ and $F = \frac{1}{2}(\varphi(x-y)+\psi(x+y), \varphi(x-y)-\psi(x+y))$ where both (α, β) and (φ, ψ) are in H^- . Then, we have $G \circ F(x, y) = \frac{1}{2}(\alpha \circ \psi(x+y) + \beta \circ \varphi(x-y), \alpha \circ \psi(x+y) - \beta \circ \varphi(x-y))$ and thus $\Pi(G \circ F) = (\alpha \circ \psi, \beta \circ \varphi)$. Since $\Pi(G) = (\alpha, \beta)$ and $\Pi(F) = (\varphi, \psi)$, we have $\Pi(G \circ F) = (\alpha \circ \psi, \beta \circ \varphi) = (\alpha, \beta) \circ \pi(\alpha, \beta) \cdot (\varphi, \psi) = \Pi(G) * \Pi(F)$.

This show that Π is an isomorphism and the proof is completed. $\square$

In Ref. [3], it is shown that $H(\mathbb{R})$ is a subgroup of G and this can also be seen in the above theorem as follows. If we define a map $\Omega : H(\mathbb{R}) \rightarrow H$ by $\Omega(f) = (f, f)$, then it is easy to see that Ω is an injective homomorphism and thus, $H(\mathbb{R})$ is a subgroup of G through an injective homomorphism $\Pi^{-1} \circ \Omega$. Zeeman's result tells us that the group of causal automorphisms on $\mathbb{R}_1^n$ is finite dimensional when $n \geq 3$ and our result tells us that the group is infinite dimensional when $n = 2$.

4. ACKNOWLEDGEMENT

This work was supported by a grant from the College of Applied Science, Kyung Hee University research professor fellowship.

REFERENCES

- [1] E.C. Zeeman, *Causality implies the Lorentz group*, J. Math. Phys. **5**, (1964) pp.490.
- [2] D.-H. Kim, *Causal automorphisms of two-dimensional Minkowski space-time*, Class. Quantum. Grav. To appear
- [3] D.-H. Kim, *An imbedding of Lorentzian manifolds*, Class. Quantum. Grav. **26**, (2009) pp. 075004.

DEPARTMENT OF APPLIED MATHEMATICS, COLLEGE OF APPLIED SCIENCE, KYUNG
HEE UNIVERSITY, SEOCHON-DONG, GIHEUNG-GU, YONGIN-SI, GYEONGGI-DO 446-701,
REPUBLIC OF KOREA

E-mail address: mathph@khu.ac.kr